



CHANCELLOR PARK
FAMILY MEDICAL PRACTICE
P.O Box 7075, Sippy Downs Q 4555
Ph: 5445 6853, Fax: 5445 2405

POST VACCINATION PARENT ADVICE SHEET

Patient Surname: _____ Patient Address: _____
Patient Given Names: _____
Patient D.O.B _____
Seen by Doctor: _____
Vaccines given by Nurse: _____

Below are listed some of the common reactions experienced following immunisation. The consent form also lists much rarer and more serious complications of which you should be aware. If symptoms other than those listed occur, or the listed symptoms persist for longer than indicated, please contact the Practice for advice.

MMR Measles, Mumps and Rubella
<ul style="list-style-type: none">Occasionally injection site nodule – may last many weeks (no treatment needed) <p>Seen 7-10 days after vaccination:</p> <ul style="list-style-type: none">Low grade temperature (fever) lasting 2-3 days, faint red rash (not infectious), headcold and/or runny nose, cough and/or puffy eyesDrowsiness or tirednessSwelling of salivary glands
Meningococcal Meningitis C
<ul style="list-style-type: none">Irritable, crying, unsettled and generally unhappyLoss of appetiteHeadache (usually observed in adolescent/adults)Localised pain, redness and swelling at injection siteOccasionally injection site nodule – may last many weeks (no treatment needed)Low grade temperature (fever)
HepB – Hib Hepatitis B, Haemophilus Influenzae B
<ul style="list-style-type: none">Localised pain, redness and swelling at injection siteOccasionally injection site nodule – may last many weeks (no treatment needed)Low grade temperature (fever)

Varilrix Chicken Pox
<ul style="list-style-type: none">Localised pain, redness and swelling at injection siteOccasional injection site nodule – may last many weeks (no treatment needed)Low grade temperature (fever) <p>Seen 5-26 days after vaccination:</p> <ul style="list-style-type: none">Pustular rash (2-5 lesions) usually at injection site which occasionally covers other parts of the body
DTPa – IPV Diphtheria, Tetanus, Whooping Cough, Polio
<ul style="list-style-type: none">Irritable, crying, unsettled and generally unhappyDrowsiness or tirednessLocalised pain, redness and swelling at injection siteOccasionally injection site nodule – may last many weeks (no treatment needed)Low grade temperature (fever)
Prevenar Pneumococcal
<ul style="list-style-type: none">Localised pain, redness and swelling at injection siteOccasionally injection site nodule – may last many weeks (no treatment needed)Low grade temperature (fever)